

Linearized Alternating Direction Method: Two Blocks and Multiple Blocks



Zhouchen Lin

林宙辰

北京大学

Dec. 23, 2014

Outline

- Alternating Direction Method (ADM)
- Linearized Alternating Direction Method (LADM) – Two Blocks
- LADM – Multiple Blocks
- Proximal LADM – Multiple Blocks
- Conclusions

Background

- Optimization is everywhere

Compressed Sensing: $\min_x \|x\|_1, \quad s.t. \quad Ax = b.$

RPCA w/ Missing Value: $\min \|A\|_* + \lambda\|E\|_1, \quad s.t. \quad \pi_\Lambda(A + E) = d.$

LASSO: $\min_x \|Ax - b\|_2, \quad s.t. \quad \|x\|_1 \leq \varepsilon.$

Image Restoration: $\min_x \|Ax - b\|_2^2 + \lambda\|\nabla x\|_1, \quad s.t. \quad 0 \leq x \leq 255.$

Covariance Selection: $\min_X \text{tr}(\Sigma X) - \log(\det(X)) + \rho e^T |X| e,$

$s.t. \quad X \in S_\Lambda^n, \text{ where } S_\Lambda^n = \{X \succcurlyeq 0 \mid \lambda_{\min} I \preccurlyeq X \preccurlyeq \lambda_{\max} I\}$

Pose Estimation: $\min_Q \text{tr}(WQ), \quad s.t. \quad \text{tr}(A_i Q) = 0, i = 1, \dots, m,$

$Q \succcurlyeq 0, \text{rank}(Q) \leq 1.$

Alternating Direction Method (ADM)

Model Problem:

$$\begin{aligned} \min_{\mathbf{x}_1, \mathbf{x}_2} \quad & f_1(\mathbf{x}_1) + f_2(\mathbf{x}_2), \\ \text{s.t.} \quad & \mathcal{A}_1(\mathbf{x}_1) + \mathcal{A}_2(\mathbf{x}_2) = \mathbf{b}, \end{aligned}$$

Augmented
Lagrangian Function



where f_i are convex functions and \mathcal{A}_i are linear mappings.

$$\begin{aligned} \tilde{\mathcal{L}}(\mathbf{x}_1, \mathbf{x}_2, \lambda) = \quad & f_1(\mathbf{x}_1) + f_2(\mathbf{x}_2) + \langle \lambda, \mathcal{A}_1(\mathbf{x}_1) + \mathcal{A}_2(\mathbf{x}_2) - \mathbf{b} \rangle \\ & + \frac{\beta}{2} \|\mathcal{A}_1(\mathbf{x}_1) + \mathcal{A}_2(\mathbf{x}_2) - \mathbf{b}\|_F^2, \end{aligned}$$

$$\mathbf{x}_1^{k+1} = \arg \min_{\mathbf{x}_1} \tilde{\mathcal{L}}(\mathbf{x}_1, \mathbf{x}_2^k, \lambda^k),$$

$$\mathbf{x}_2^{k+1} = \arg \min_{\mathbf{x}_2} \tilde{\mathcal{L}}(\mathbf{x}_1^{k+1}, \mathbf{x}_2, \lambda^k),$$

$$\lambda^{k+1} = \lambda^k + \beta_k [\mathcal{A}_1(\mathbf{x}_1^{k+1}) + \mathcal{A}_2(\mathbf{x}_2^{k+1}) - \mathbf{b}].$$

Assume: Easy



Update β_k

Linearized Alternating Direction Method (LADM)

$$\mathbf{x}_1^{k+1} = \arg \min_{\mathbf{x}_1} f_1(\mathbf{x}_1) + \frac{\beta_k}{2} \|\mathcal{A}_1(\mathbf{x}_1) + \mathcal{A}_2(\mathbf{x}_2^k) - \mathbf{b} + \lambda_k / \beta_k\|^2,$$

$$\mathbf{x}_2^{k+1} = \arg \min_{\mathbf{x}_2} f_2(\mathbf{x}_2) + \frac{\beta_k}{2} \|\mathcal{A}_2(\mathbf{x}_1^{k+1}) + \mathcal{A}_2(\mathbf{x}_2) - \mathbf{b} + \lambda_k / \beta_k\|^2$$

$$\min_{\mathbf{x}} f_1(\mathbf{x}) + \frac{\beta}{2} \|\mathbf{x} - \mathbf{w}\|_F^2$$


$$\min_{\mathbf{x}} f_2(\mathbf{x}) + \frac{\beta}{2} \|\mathbf{x} - \mathbf{w}\|_F^2$$

Proximal
Operation

$$\arg \min_{\mathbf{x}} \|\mathbf{x}\|_1 + \frac{\beta}{2} \|\mathbf{x} - \mathbf{w}\|^2 = \mathcal{T}_{\beta^{-1}}(\mathbf{w}),$$

$$\mathcal{T}_{\varepsilon}(x) = \text{sgn}(x) \max(|x| - \varepsilon, 0).$$

$$\arg \min_X \|X\|_* + \frac{\varepsilon}{2} \|X - W\|_F^2 = \Theta_{\varepsilon^{-1}}(W) = U \mathcal{T}_{\varepsilon^{-1}}(S) V^T,$$



where $W = USV^T$ is the singular value decomposition (SVD) of W .

Linearized Alternating Direction Method (LADM)

Introducing auxiliary variables:

$$\begin{aligned} \min_{\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4} \quad & f_1(\mathbf{x}_1) + f_2(\mathbf{x}_2), \\ \text{s.t.} \quad & \mathbf{x}_1 = \mathbf{x}_3, \mathbf{x}_2 = \mathbf{x}_4, \mathcal{A}_1(\mathbf{x}_3) + \mathcal{A}_2(\mathbf{x}_4) = \mathbf{b}. \end{aligned}$$

$$\begin{aligned} & \tilde{\mathcal{L}}(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4, \lambda_1, \lambda_2, \lambda_3) \\ = & f_1(\mathbf{x}_1) + f_2(\mathbf{x}_2) + \langle \lambda_1, \mathbf{x}_1 - \mathbf{x}_3 \rangle + \langle \lambda_2, \mathbf{x}_2 - \mathbf{x}_4 \rangle + \langle \lambda_3, \mathcal{A}_1(\mathbf{x}_3) + \mathcal{A}_2(\mathbf{x}_4) - \mathbf{b} \rangle \\ & + \frac{\beta}{2} (\|\mathbf{x}_1 - \mathbf{x}_3\|_F^2 + \|\mathbf{x}_2 - \mathbf{x}_4\|_F^2 + \|\mathcal{A}_1(\mathbf{x}_3) + \mathcal{A}_2(\mathbf{x}_4) - \mathbf{b}\|_F^2), \end{aligned}$$

Three drawbacks:

1. More blocks \longrightarrow more memory & slower convergence.
2. Matrix inversion is expensive.
3. Convergence is NOT guaranteed!

Linearized Alternating Direction Method (LADM)

$$\begin{aligned}\mathbf{x}_1^{k+1} &= \arg \min_{\mathbf{x}_1} f_1(\mathbf{x}_1) + \frac{\beta_k}{2} \|\mathcal{A}_1(\mathbf{x}_1) + \mathcal{A}_2(\mathbf{x}_2^k) - \mathbf{b} + \lambda_k / \beta_k\|^2, \\ \mathbf{x}_2^{k+1} &= \arg \min_{\mathbf{x}_2} f_2(\mathbf{x}_2) + \frac{\beta_k}{2} \|\mathcal{A}_2(\mathbf{x}_1^{k+1}) + \mathcal{A}_2(\mathbf{x}_2) - \mathbf{b} + \lambda_k / \beta_k\|^2\end{aligned}$$

$$\min_{\mathbf{x}} f_1(\mathbf{x}) + \frac{\beta}{2} \|\mathbf{x} - \mathbf{w}\|_F^2$$

$$\min_{\mathbf{x}} f_2(\mathbf{x}) + \frac{\beta}{2} \|\mathbf{x} - \mathbf{w}\|_F^2$$

- Linearize the quadratic term

$$\begin{aligned}\mathbf{x}_1^{k+1} &= \arg \min_{\mathbf{x}_1} f_1(\mathbf{x}_1) + \langle \mathcal{A}_1^*(\lambda_k) + \beta_k \mathcal{A}_1^*(\mathcal{A}_1(\mathbf{x}_1^k) + \mathcal{A}_2(\mathbf{x}_2^k) - \mathbf{b}), \mathbf{x}_1 - \mathbf{x}_1^k \rangle \\ &\quad + \frac{\beta_k \eta_1}{2} \|\mathbf{x}_1 - \mathbf{x}_1^k\|^2 \\ &= \arg \min_{\mathbf{x}_1} f_1(\mathbf{x}_1) \\ &\quad + \frac{\beta_k \eta_1}{2} \|\mathbf{x}_1 - \mathbf{x}_1^k + \mathcal{A}_1^*(\lambda_k + \beta_k (\mathcal{A}_1(\mathbf{x}_1^k) + \mathcal{A}_2(\mathbf{x}_2^k) - \mathbf{b})) / (\beta_k \eta_1)\|^2,\end{aligned}$$

$$\langle \mathcal{A}^*(x), y \rangle = \langle x, \mathcal{A}(y) \rangle, \quad \forall x, y$$

Adjoint Operator

$$\begin{aligned}\mathbf{x}_2^{k+1} &= \arg \min_{\mathbf{x}_2} f_2(\mathbf{x}_2) \\ &\quad + \frac{\beta_k \eta_2}{2} \|\mathbf{x}_2 - \mathbf{x}_2^k + \mathcal{A}_2^*(\lambda_k + \beta_k (\mathcal{A}_1(\mathbf{x}_1^{k+1}) + \mathcal{A}_2(\mathbf{x}_2^k) - \mathbf{b})) / (\beta_k \eta_2)\|^2.\end{aligned}$$

LADM with Adaptive Penalty (LADMAP)

Theorem: If $\{\beta_k\}$ is non-decreasing and upper bounded, $\eta_i > \|\mathcal{A}_i\|^2$, $i = 1, 2$, then the sequence $\{(\mathbf{x}_1^k, \mathbf{x}_2^k, \lambda_k)\}$ converges to a KKT point of the model problem.

LADM with Adaptive Penalty (LADMAP)

- Adaptive Penalty

$$\begin{aligned} \mathbf{x}_1^{k+1} &= \arg \min_{\mathbf{x}_1} f_1(\mathbf{x}_1) \\ &\quad + \frac{\beta_k \eta_1}{2} \|\mathbf{x}_1 - \mathbf{x}_1^k + \mathcal{A}_1^*(\lambda_k + \beta_k(\mathcal{A}_1(\mathbf{x}_1^k) + \mathcal{A}_2(\mathbf{x}_1^k) - \mathbf{b})) / (\beta_k \eta_1)\|^2, \end{aligned}$$

$$\begin{aligned} \mathbf{x}_2^{k+1} &= \arg \min_{\mathbf{x}_2} f_2(\mathbf{x}_2) \\ &\quad + \frac{\beta_k \eta_2}{2} \|\mathbf{x}_2 - \mathbf{x}_2^k + \mathcal{A}_2^*(\lambda_k + \beta_k(\mathcal{A}_1(\mathbf{x}_1^{k+1}) + \mathcal{A}_2(\mathbf{x}_2^k) - \mathbf{b})) / (\beta_k \eta_2)\|^2. \end{aligned}$$

⇓

$$\begin{aligned} -\beta_k \eta_1 (\mathbf{x}_1^{k+1} - \mathbf{x}_1^k) - \mathcal{A}_1^*(\lambda_k + \beta_k(\mathcal{A}_1(\mathbf{x}_1^k) + \mathcal{A}_2(\mathbf{x}_1^k) - \mathbf{b})) &\in \partial f_1(\mathbf{x}_1^{k+1}) \\ -\beta_k \eta_2 (\mathbf{x}_2^{k+1} - \mathbf{x}_2^k) - \mathcal{A}_2^*(\lambda_k + \beta_k(\mathcal{A}_1(\mathbf{x}_1^{k+1}) + \mathcal{A}_2(\mathbf{x}_2^k) - \mathbf{b})) &\in \partial f_2(\mathbf{x}_2^{k+1}) \end{aligned}$$

KKT condition: $\exists(\mathbf{x}^*, \mathbf{y}^*, \lambda^*)$ such that

$$\begin{aligned} \mathcal{A}_1(\mathbf{x}_1^*) + \mathcal{A}_2(\mathbf{x}_2^*) - \mathbf{b} &= \mathbf{0}, \\ -\mathcal{A}_1^*(\lambda^*) &\in \partial f_1(\mathbf{x}_1^*), \quad -\mathcal{A}_2^*(\lambda^*) \in \partial f_2(\mathbf{x}_2^*). \end{aligned}$$

LADM with Adaptive Penalty (LADMAP)

Both $\beta_k \eta_1 \|\mathbf{x}_1^{k+1} - \mathbf{x}_1^k\| / \|\mathcal{A}_1^*(\mathbf{b})\|$ and $\beta_k \eta_2 \|\mathbf{x}_2^{k+1} - \mathbf{x}_2^k\| / \|\mathcal{A}_2^*(\mathbf{b})\|$ should be small.

$$\eta_i = \|\mathcal{A}_i\|^2 \quad \Rightarrow \quad \text{Approximate } \|\mathcal{A}_i^*(\mathbf{b})\| \text{ by } \sqrt{\eta_i} \|\mathbf{b}\|$$

- Adaptive Penalty

$$\beta_{k+1} = \min(\beta_{\max}, \rho \beta_k),$$

$$\rho = \begin{cases} \rho_0, & \text{if } \beta_k \max(\sqrt{\eta_1} \|\mathbf{x}_1^{k+1} - \mathbf{x}_1^k\|_F, \sqrt{\eta_2} \|\mathbf{x}_2^{k+1} - \mathbf{x}_2^k\|_F) / \|\mathbf{b}\|_F < \varepsilon_2, \\ 1, & \text{otherwise,} \end{cases}$$

where $\rho_0 \geq 1$ is a constant.

- Loop until


$$\|\mathcal{A}_1(\mathbf{x}_1^{k+1}) + \mathcal{A}_2(\mathbf{x}_2^{k+1}) - \mathbf{b}\|_F < \varepsilon_1,$$

$$\beta_k \max(\sqrt{\eta_1} \|\mathbf{x}_1^{k+1} - \mathbf{x}_1^k\|_F, \sqrt{\eta_2} \|\mathbf{x}_2^{k+1} - \mathbf{x}_2^k\|_F) / \|\mathbf{b}\|_F < \varepsilon_2.$$

LADM with Adaptive Penalty (LADMAP)

- Choice of parameters

1. $\beta_0 = \alpha \varepsilon_2$, where $\alpha \propto$ the size of \mathbf{b} . β_0 should not be too large, so that β_k increases in the first few iterations.
2. $\rho_0 \geq 1$ should be chosen such that β_k increases steadily (but not necessarily every iteration).



LADM with Adaptive Penalty (LADM-AP)

- An example (LRR):

$$\min_{Z, E} \|Z\|_* + \mu \|E\|_1, \quad s.t. \quad X = XZ + E.$$

$$\mathcal{A}_1(Z) = XZ, \quad \mathcal{A}_2(E) = E.$$

$$\mathcal{A}_1^*(Z) = X^T Z, \quad \mathcal{A}_2^*(E) = E, \quad \eta_1 = \|X\|_2^2, \quad \eta_2 = 1.$$

Experiment

Table 1: Comparison among APG, ADM, LADM and LADMAP on the synthetic data. For each quadruple (s, p, d, \tilde{r}) , the LRR problem, with $\mu = 0.1$, was solved for the same data using different algorithms. We present typical running time (in $\times 10^3$ seconds), iteration number, relative error (%) of output solution $(\hat{\mathbf{E}}, \hat{\mathbf{Z}})$ and the clustering accuracy (%) of tested algorithms, respectively.

Size (s, p, d, \tilde{r})	Method	Time	Iter.	$\frac{\ \tilde{\mathbf{Z}} - \mathbf{Z}_0\ }{\ \mathbf{Z}_0\ }$	$\frac{\ \hat{\mathbf{E}} - \mathbf{E}_0\ }{\ \mathbf{E}_0\ }$	Acc.
(10, 20,200, 5)	APG	0.0332	110	2.2079	1.5096	81.5
	ADM	0.0529	176	0.5491	0.5093	90.0
	LADM	0.0603	194	0.5480	0.5024	90.0
	LADMAP	0.0145	46	0.5480	0.5024	90.0
(15, 20,300, 5)	APG	0.0869	106	2.4824	1.0341	80.0
	ADM	0.1526	185	0.6519	0.4078	83.7
	LADM	0.2943	363	0.6518	0.4076	86.7
	LADMAP	0.0336	41	0.6518	0.4076	86.7
(20, 25, 500, 5)	APG	1.8837	117	2.8905	2.4017	72.4
	ADM	3.7139	225	1.1191	1.0170	80.0
	LADM	8.1574	508	0.6379	0.4268	80.0
	LADMAP	0.7762	40	0.6379	0.4268	84.6
(30, 30, 900, 5)	APG	6.1252	116	3.0667	0.9199	69.4
	ADM	11.7185	220	0.6865	0.4866	76.0
	LADM	N.A.	N.A.	N.A.	N.A.	N.A.
	LADMAP	2.3891	44	0.6864	0.4294	80.1

LADM with Parallel Splitting and Adaptive Penalty (LADMPSAP)

- Model problem:

$$\min_{\mathbf{x}_1, \dots, \mathbf{x}_n} \sum_{i=1}^n f_i(\mathbf{x}_i), \quad s.t. \quad \sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i) = \mathbf{b}.$$

$$\min_{\mathbf{X}} \|\mathbf{X}\|_* + \frac{1}{2\mu} \|\mathbf{b} - \mathcal{P}(\mathbf{X})\|^2, \quad s.t. \quad \mathbf{X} \geq 0,$$

⇓

$$\min_{\mathbf{X}, \mathbf{e}} \|\mathbf{X}\|_* + \frac{1}{2\mu} \|\mathbf{e}\|^2, \quad s.t. \quad \mathbf{b} = \mathcal{P}(\mathbf{X}) + \mathbf{e}, \quad \mathbf{X} \geq 0,$$

⇓

$$\min_{\mathbf{X}, \mathbf{Y}, \mathbf{e}} \|\mathbf{X}\|_* + \frac{1}{2\mu} \|\mathbf{e}\|^2, \quad s.t. \quad \mathbf{b} = \mathcal{P}(\mathbf{Y}) + \mathbf{e}, \quad \mathbf{X} = \mathbf{Y}, \quad \mathbf{Y} \geq 0,$$

⇓

$$\min_{\mathbf{X}, \mathbf{Y}, \mathbf{e}} \|\mathbf{X}\|_* + \frac{1}{2\mu} \|\mathbf{e}\|^2 + \chi_{\mathbf{Y} \geq 0}(\mathbf{Y}), \quad s.t. \quad \mathbf{b} = \mathcal{P}(\mathbf{Y}) + \mathbf{e}, \quad \mathbf{X} = \mathbf{Y}.$$

LADM with Parallel Splitting and Adaptive Penalty (LADMPSAP)

- Can we naively generalize two-block LADMAP for multi-block problems?

No!

Actually, the naive generalization of LADMAP may be divergent, e.g., when applied to the following problem with $n \geq 5$:

$$\min_{\mathbf{x}_1, \dots, \mathbf{x}_n} \sum_{i=1}^n \|\mathbf{x}_i\|_1, \quad s.t. \quad \sum_{i=1}^n \mathbf{A}_i \mathbf{x}_i = \mathbf{b}.$$

Lin et al., *Linearized Alternating Direction Method with Parallel Splitting and Adaptive Penalty for Separable Convex Programs in Machine Learning*, ML, 2015.

C. Chen et al. *The Direct Extension of ADMM for Multi-block Convex Minimization Problems is Not Necessarily Convergent*. Preprint.

LADM with Parallel Splitting and Adaptive Penalty (LADMPSAP)

$$\mathbf{x}_i^{k+1} = \operatorname{argmin}_{\mathbf{x}_i} f_i(\mathbf{x}_i) + \frac{\eta_i \beta_k}{2} \left\| \mathbf{x}_i - \mathbf{x}_i^k + \mathcal{A}_i^* \left(\lambda^k + \beta_k \left(\sum_{j=1}^n \mathcal{A}_i(\mathbf{x}_j^k) - \mathbf{b} \right) \right) / (\eta_i \beta_k) \right\|^2,$$

$$i = 1, \dots, n,$$

$$\sum_{j=1}^{i-1} \mathcal{A}_i(\mathbf{x}_j^{k+1}) + \sum_{j=i}^n \mathcal{A}_i(\mathbf{x}_j^k)$$

$$\lambda^{k+1} = \lambda^k + \beta_k \left(\sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i^{k+1}) - \mathbf{b} \right),$$

Parallel!

$$\beta_{k+1} = \min(\beta_{\max}, \rho \beta_k),$$

where

$$\rho = \begin{cases} \rho_0, & \text{if } \beta_k \max(\{\sqrt{\eta_i} \|\mathbf{x}_i^{k+1} - \mathbf{x}_i^k\|, i = 1, \dots, n\}) / \|\mathbf{b}\| < \varepsilon_2, \\ 1, & \text{otherwise,} \end{cases}$$

with $\rho_0 > 1$ being a constant and $0 < \varepsilon_2 \ll 1$ being a threshold.

LADM with Parallel Splitting and Adaptive Penalty (LADMPSAP)

Theorem: If $\{\beta_k\}$ is non-decreasing and upper bounded, $\eta_i > n\|A_i\|^2$, $i = 1, \dots, n$, then $\{(\{\mathbf{x}_i^k\}, \lambda^k)\}$ generated by LADMPSAP converges to a KKT point of the problem.

Remark: When $n = 2$, LADMPSAP is weaker than LADMAP:

$$\eta_i > 2\|A_i\|^2 \text{ vs. } \eta_i > \|A_i\|^2.$$

- Related work: He & Yuan, *Linearized Alternating Direction Method with Gaussian Back Substitution for Separable Convex Programming*, preprint.

LADM with Parallel Splitting and Adaptive Penalty (LADMPSAP)

- Model problem:

$$\min_{\mathbf{x}_1, \dots, \mathbf{x}_n} \sum_{i=1}^n f_i(\mathbf{x}_i), \quad s.t. \quad \sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i) = \mathbf{b}, \quad \mathbf{x}_i \in X_i, i = 1, \dots, n,$$

where $X_i \subseteq \mathbb{R}^{d_i}$ is a closed convex set.

⇓

$$\min_{\mathbf{x}_1, \dots, \mathbf{x}_{2n}} \sum_{i=1}^n f_i(\mathbf{x}_i) + \sum_{i=n+1}^{2n} \chi_{\mathbf{x}_i \in X_{i-n}}(\mathbf{x}_i), \quad s.t. \quad \sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i) = \mathbf{b}, \quad \mathbf{x}_i = \mathbf{x}_{n+i}, i = 1, \dots, n.$$

Theorem: If $\{\beta_k\}$ is non-decreasing and upper bounded, $\mathbf{x}_{n+1}, \dots, \mathbf{x}_{2n}$ are auxiliary variables, $\eta_i > n\|\mathcal{A}_i\|^2 + 2, \eta_{n+i} > 2, i = 1, \dots, n$, then $\{(\{\mathbf{x}_i^k\}, \lambda^k)\}$ generated by LADMPSAP converges to a KKT point of the problem.

$$\eta_i > 2n(\|\mathcal{A}_i\|^2 + 1), \eta_{n+i} > 2n, i = 1, \dots, n$$

Experiment

$$\min_{\mathbf{X}} \|\mathbf{X}\|_* + \frac{1}{2\mu} \|\mathbf{b} - \mathcal{P}(\mathbf{X})\|^2, \quad s.t. \quad \mathbf{X} \geq 0,$$

⇓

$$\min_{\mathbf{X}, \mathbf{Y}, \mathbf{e}} \|\mathbf{X}\|_* + \frac{1}{2\mu} \|\mathbf{e}\|^2 + \chi_{\mathbf{Y} \geq 0}(\mathbf{Y}), \quad s.t. \quad \mathbf{b} = \mathcal{P}(\mathbf{Y}) + \mathbf{e}, \quad \mathbf{X} = \mathbf{Y}.$$



(a) Original

(b) Corrupted

(c) FPCA

(d) LADM

(e) LADMPSAP

Experiment

Table 1: Numerical comparison on the NMC problem with synthetic data, average of 10 runs. q , t and d_r denote, respectively, sample ratio, the number of measurements $t = q(mn)$ and the “degree of freedom” defined by $d_r = r(m + n - r)$ for a matrix with rank r and q . Here we set $m = n$ and fix $r = 10$ in all the tests.

X			LADM				LADMPSAP			
n	q	t/d_r	Iter.	Time(s)	RelErr	FA	Iter.	Time(s)	RelErr	FA
1000	20%	10.05	375	177.92	1.35E-5	6.21E-4	58	24.94	9.67E-6	0
	10%	5.03	1000	459.70	4.60E-5	6.50E-4	109	42.68	1.72E-5	0
5000	20%	50.05	229	1613.68	1.08E-5	1.93E-4	49	369.96	9.05E-6	0
	10%	25.03	539	2028.14	1.20E-5	7.70E-5	89	365.26	9.76E-6	0
10000	10%	50.03	463	6679.59	1.11E-5	4.18E-5	89	1584.39	1.03E-5	0

Table 1: Numerical comparison on the image inpainting problem.

Method	#Iter.	Time(s)	PSNR	FA
FPCA	179	228.99	27.77dB	9.41E-4
LADM	228	207.95	26.98dB	2.92E-3
LADMPSAP	143	134.89	31.39dB	0

LADM with Parallel Splitting and Adaptive Penalty (LADMPSAP)

Enhanced convergence results:

Theorem 1: If $\{\beta_k\}$ is non-decreasing and $\sum_{k=1}^{+\infty} \beta_k^{-1} = +\infty$, $\eta_i > n\|\mathcal{A}_i\|^2$, $\partial f_i(\mathbf{x})$ is bounded, $i = 1, \dots, n$, then the sequence $\{\mathbf{x}_i^k\}$ generated by LADMPSAP converges to an optimal solution to the model problem.

Theorem 2: If $\{\beta_k\}$ is non-decreasing, $\eta_i > n\|\mathcal{A}_i\|^2$, $\partial f_i(\mathbf{x})$ is bounded, $i = 1, \dots, n$, then $\sum_{k=1}^{+\infty} \beta_k^{-1} = +\infty$ is also the necessary condition for the global convergence of $\{\mathbf{x}_i^k\}$ generated by LADMPSAP to an optimal solution to the model problem.

With the above analysis, when all the subgradients of the component objective functions are bounded we can remove the upper bound β_{\max} .

LADM with Parallel Splitting and Adaptive Penalty (LADMPSAP)

Define $\mathbf{x} = (\mathbf{x}_1^T, \dots, \mathbf{x}_n^T)^T$, $\mathbf{x}^* = ((\mathbf{x}_1^*)^T, \dots, (\mathbf{x}_n^*)^T)^T$ and $f(\mathbf{x}) = \sum_{i=1}^n f_i(\mathbf{x}_i)$, where $(\mathbf{x}_1^*, \dots, \mathbf{x}_n^*, \lambda^*)$ is a KKT point of the model problem.

Proposition: $\tilde{\mathbf{x}}$ is an optimal solution to the model problem iff there exists $\alpha > 0$, such that

$$f(\tilde{\mathbf{x}}) - f(\mathbf{x}^*) + \sum_{i=1}^n \langle \mathcal{A}_i^*(\lambda^*), \tilde{\mathbf{x}}_i - \mathbf{x}_i^* \rangle + \alpha \left\| \sum_{i=1}^n \mathcal{A}_i(\tilde{\mathbf{x}}_i) - \mathbf{b} \right\|^2 = 0.$$

Our criterion for checking the optimality of a solution is much simpler than that in He et al. 2011, which has to compare with all $(\mathbf{x}_1, \dots, \mathbf{x}_n, \lambda) \in \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_n} \times \mathbb{R}^m$.

LADM with Parallel Splitting and Adaptive Penalty (LADMPSAP)

Theorem 3: Define $\bar{\mathbf{x}}^K = \sum_{k=0}^K \gamma_k \mathbf{x}^{k+1}$, where $\gamma_k = \beta_k^{-1} / \sum_{j=0}^K \beta_j^{-1}$. Then

$$f(\bar{\mathbf{x}}^K) - f(\mathbf{x}^*) + \sum_{i=1}^n \langle \mathcal{A}_i^*(\lambda^*), \bar{\mathbf{x}}_i^K - \mathbf{x}_i^* \rangle + \frac{\alpha\beta_0}{2} \left\| \sum_{i=1}^n \mathcal{A}_i(\bar{\mathbf{x}}_i^K) - \mathbf{b} \right\|^2 \leq C_0 / \left(2 \sum_{k=0}^K \beta_k^{-1} \right), \quad (1)$$

where $\alpha^{-1} = (n+1) \max \left(1, \left\{ \frac{\|\mathcal{A}_i\|^2}{\eta_i - n\|\mathcal{A}_i\|^2}, i = 1, \dots, n \right\} \right)$ and $C_0 = \sum_{i=1}^n \eta_i \|\mathbf{x}_i^0 - \mathbf{x}_i^*\|^2 + \beta_0^{-2} \|\lambda^0 - \lambda^*\|^2$.

A much simpler proof of convergence rate (in ergodic sense)!

Proximal LADMPSAP

- Even more general problem:

$$\min_{\mathbf{x}_1, \dots, \mathbf{x}_n} \sum_{i=1}^n f_i(\mathbf{x}_i), \quad s.t. \quad \sum_{i=1}^n \mathcal{A}_i(\mathbf{x}_i) = \mathbf{b}.$$

$$f_i(\mathbf{x}_i) = g_i(\mathbf{x}_i) + h_i(\mathbf{x}_i),$$

where both g_i and h_i are convex, g_i is $C^{1,1}$:

$$\|\nabla g_i(\mathbf{x}) - \nabla g_i(\mathbf{y})\| \leq L_i \|\mathbf{x} - \mathbf{y}\|, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^{d_i},$$

and h_i may not be differentiable but its proximal operation is easily solvable.

Proximal LADMPSAP

- Linearize the augmented term to obtain:

$$\mathbf{x}_i^{k+1} = \underset{\mathbf{x}_i}{\operatorname{argmin}} h_i(\mathbf{x}_i) + g_i(\mathbf{x}_i) + \frac{\sigma_i^{(k)}}{2} \left\| \mathbf{x}_i - \mathbf{x}_i^k + \mathcal{A}_i^\dagger(\hat{\lambda}^k)/\sigma_i^{(k)} \right\|^2, \quad i = 1, \dots, n,$$

- Further linearize g_i :

$$\begin{aligned} \mathbf{x}_i^{k+1} &= \underset{\mathbf{x}_i}{\operatorname{argmin}} h_i(\mathbf{x}_i) + g_i(\mathbf{x}_i^k) + \frac{\sigma_i^{(k)}}{2} \left\| \mathcal{A}_i^\dagger(\hat{\lambda}^k)/\sigma_i^{(k)} \right\|^2 \\ &\quad + \langle \nabla g_i(\mathbf{x}_i^k) + \mathcal{A}_i^\dagger(\hat{\lambda}^k), \mathbf{x}_i - \mathbf{x}_i^k \rangle + \frac{\tau_i^{(k)}}{2} \left\| \mathbf{x}_i - \mathbf{x}_i^k \right\|^2 \\ &= \underset{\mathbf{x}_i}{\operatorname{argmin}} h_i(\mathbf{x}_i) + \frac{\tau_i^{(k)}}{2} \left\| \mathbf{x}_i - \mathbf{x}_i^k + \frac{1}{\tau_i^{(k)}} [\mathcal{A}_i^\dagger(\hat{\lambda}^k) + \nabla g_i(\mathbf{x}_i^k)] \right\|^2. \end{aligned}$$

- Convergence condition:

$$\tau_i^{(k)} = T_i + \beta_k \eta_i, \text{ where } T_i \geq L_i \text{ and } \eta_i > n \|\mathcal{A}_i\|^2 \text{ are both positive constants.}$$

Experiment

- Group Sparse Logistic Regression with Overlap

$$\min_{\mathbf{w}, b} \frac{1}{s} \sum_{i=1}^s \log (1 + \exp (-y_i(\mathbf{w}^T \mathbf{x}_i + b))) + \mu \sum_{j=1}^t \|\mathbf{S}_j \mathbf{w}\|, \quad (1)$$

where \mathbf{x}_i and y_i , $i = 1, \dots, s$, are the training data and labels, respectively, and \mathbf{w} and b parameterize the linear classifier. \mathbf{S}_j , $j = 1, \dots, t$, are the selection matrices, with only one 1 at each row and the rest entries are all zeros. The groups of entries, $\mathbf{S}_j \mathbf{w}$, $j = 1, \dots, t$, may overlap each other.

Introducing $\bar{\mathbf{w}} = (\mathbf{w}^T, b)^T$, $\bar{\mathbf{x}}_i = (\mathbf{x}_i^T, 1)^T$, $\mathbf{z} = (\mathbf{z}_1^T, \mathbf{z}_2^T, \dots, \mathbf{z}_t^T)^T$, and $\bar{\mathbf{S}} = (\mathbf{S}, \mathbf{0})$, where $\mathbf{S} = (\mathbf{S}_1^T, \dots, \mathbf{S}_t^T)^T$, (1) can be rewritten as

$$\min_{\bar{\mathbf{w}}, \mathbf{z}} \frac{1}{s} \sum_{i=1}^s \log (1 + \exp (-y_i(\bar{\mathbf{w}}^T \bar{\mathbf{x}}_i))) + \mu \sum_{j=1}^t \|\mathbf{z}_j\|, \quad s.t. \quad \mathbf{z} = \bar{\mathbf{S}} \bar{\mathbf{w}}, \quad (2)$$

The Lipschitz constant of the gradient of logistic function with respect to $\bar{\mathbf{w}}$ can be proven to be $L_{\bar{\mathbf{w}}} \cdot \frac{1}{4s} \|\bar{\mathbf{X}}\|_2^2$, where $\bar{\mathbf{X}} = (\bar{\mathbf{x}}_1, \bar{\mathbf{x}}_2, \dots, \bar{\mathbf{x}}_s)$.

Experiment

(s, p, t, q)	Method	Time	#Iter.	$\frac{\ \hat{\mathbf{w}} - \bar{\mathbf{w}}^*\ }{\ \bar{\mathbf{w}}^*\ }$	$\frac{\ \hat{\mathbf{z}} - \mathbf{z}^*\ }{\ \mathbf{z}^*\ }$
(300, 901, 100, 10)	ADM	294.15	43	0.4800	0.4790
	LADM	229.03	43	0.5331	0.5320
	LADMPS	105.50	47	0.2088	0.2094
	LADMPSAP	57.46	39	0.0371	0.0368
	pLADMPSAP	1.97	141	0.0112	0.0112
(450, 1351, 150, 15)	ADM	450.96	33	0.4337	0.4343
	LADM	437.12	36	0.5126	0.5133
	LADMPS	201.30	39	0.1938	0.1937
	LADMPSAP	136.64	37	0.0321	0.0306
	pLADMPSAP	4.16	150	0.0131	0.0131
(600, 1801, 200, 20)	ADM	1617.09	62	1.4299	1.4365
	LADM	1486.23	63	1.5200	1.5279
	LADMPS	494.52	46	0.4915	0.4936
	LADMPSAP	216.45	32	0.0787	0.0783
	pLADMPSAP	5.77	127	0.0276	0.0277

Conclusions

- LADMAP, LADMPSAP, and P-LADMPSAP are very general methods for solving various convex programs.
- Adaptive penalty is important for fast convergence.

Thanks!

- zlin@pku.edu.cn
 - <http://www.cis.pku.edu.cn/faculty/vision/zlin/zlin.htm>

